Precision RENORM diffraction Predictions at LHC: A puzzle solving tool?



Konstantin Goulianos
http://physics.rockefeller.edu/dino/my.html

What puzzle?

CONTENTS

- Diffraction
  - SD1 $p_1p_2 \rightarrow p_1+gap+X_2$ Single Diffraction / Dissociation –1
  - SD2 $p_1p_2 \rightarrow X_1+gap+p_2$ Single Diffraction / Dissociation - 2
  - DD $p_1p_2 \rightarrow X_1+gap+X_2$ Double Diffraction / Double Dissociation
  - CD/DPE $p_1p_2 \rightarrow gap+X+gap$ Central Diffraction / Double Pomeron Exchange

- Renormalization→Unitarization
  - **RENORM Model**
- **Triple-Pomeron Coupling: unambiguously determined** ← g_{PPP}
- **Total Cross Section**: ← σ_{tot}
  - **Unique prediction, based on a spin-2 tensor glue-ball model**
- **References**


- MBR MC Simulation in PYTHIA8, KG & R. Ciesielski, **http://arxiv.org/abs/1205.1446**

Special thanks to Robert Ciesielski, my collaborator in the PYTHIA8-MBR project

- EDS BLOIS 2015 Borgo, Corsica, France Jun 29-Jul 4, https://indico.cern.ch/event/362991/
  KG, Updated RENORM/MBR-model Predictions for Diffraction at the LHC, http://dx.doi.org/10.5506/APhysPolBSupp.8.783
- Moriond QCD 2016, La Thuile, Italy, March 19-26, http://moriond.in2p3.fr/QCD/2016/
- NPQCD16, Paris, June, https://www.brown.edu/conference/14th-workshop-non-perturbative-quantum-chromodynamics/
- **DIFFRACTION 2016**, Catania, Sep.2-8 2016 https://agenda.infn.it/conferenceDisplay.py?confId=10935 ← similar talk


RENORM: Basic and Combined Diffractive Processes

	acronym	basic diffractive processes
BASIC	$\mathbf{SD}_{\bar{p}}$	$\bar{p}p \to \bar{p} + \text{gap} + [p \to X_p]$,
	$\mathbf{SD}_{p}$	$\bar{p}p \to [\bar{p} \to X_{\bar{p}}] + \text{gap} + p$,
	$\mathbf{DD}$	$\bar{p}p \to [\bar{p} \to X_{\bar{p}}] + \text{gap} + [p \to X_p]$,
	$\mathbf{DPE}$	$\bar{p}p \to \bar{p} + \text{gap} + X_c + \text{gap} + p$,
COMBINED		2-gap combinations of SD and DD
	$\mathbf{SDD}_{\bar{p}}$	$\bar{p}p \to \bar{p} + \text{gap} + X_c + \text{gap} + [p \to X_p]$,
	$\mathbf{SDD}_{p}$	$\bar{p}p \to [\bar{p} \to X_{\bar{p}}]\text{gap} + X_c + \text{gap} + p$.



Cross sections analytically expressed in arXiv::


4-gap diffractive processes-Snowmass 2001 http://arxiv.org/abs/hep-ph/0110240



Regge Theory: Values of s_0 & g_{PPP}?

KG-PLB 358, 379 (1995)

http://www.sciencedirect.com/science/article/pii/037026939501023J



Parameters:

- s_0, s_0' and g(t)
- set $s_0' = s_0$ (universal *Pomeron*)
- determine s_0 and g_{PPP} – **how?**

$\alpha(t)=\alpha(0)+\alpha' t \quad \alpha(0)=1+\varepsilon$

$$\sigma_T = \beta_1(0)\beta_2(0)\left(\frac{s}{s_0}\right)^{\alpha(0)-1} = \sigma_0^{p\bar{p}}\left(\frac{s}{s_0}\right)^{\epsilon} \qquad (1)$$

$$\frac{d\sigma_{el}}{dt} = \frac{\beta_1^2(t)\beta_2^2(t)}{16\pi}\left(\frac{s}{s_0}\right)^{2[\alpha(t)-1]}$$

$$= \frac{\sigma_T^2}{16\pi}\left(\frac{s}{s_0}\right)^{2\alpha' t} \quad F^4(t) \approx \frac{\sigma_T^2}{16\pi} e^{b_{el}(s)t} \qquad (2)$$

$$F^4(t) \approx e^{b_{0,el}t} \quad \Rightarrow \quad b_{el}(s) = b_{0,el} + 2\alpha' \ln\left(\frac{s}{s_0}\right) \qquad (3)$$

$$\frac{d^2\sigma_{sd}}{dtd\xi}$$

$$= \frac{\beta_1^2(t)}{16\pi}\,\xi^{1-2\alpha(t)}\left[\beta_2(0)\,g(t)\left(\frac{s'}{s_0'}\right)^{\alpha(0)-1}\right]$$

$$= f_{\mathcal{P}/p}(\xi,t)\;\sigma_T^{\mathcal{P}\bar{p}}(s',t) \qquad (4)$$

Theoretical Complication: Unitarity!

$$\left(\frac{d\sigma_{el}}{dt}\right)_{t=0} \sim \left(\frac{s}{s_o}\right)^{2\epsilon}, \ \sigma_t \sim \left(\frac{s}{s_o}\right)^{\epsilon}, \ \text{and} \ \sigma_{sd} \sim \left(\frac{s}{s_o}\right)^{2\epsilon}$$

- σ_{sd} grows faster than σ_t as s increases *
  ➔ unitarity violation at high s

(also true for partial x-sections in impact parameter space)


- the unitarity limit is already reached at $\sqrt{s} \sim 2$ TeV


- need unitarization

* similarly for $(d\sigma_{el}/dt)_{t=0}$ w.r.t. σ_t, but this is handled differently in RENORM

FACTORIZATION BREAKING IN SOFT DIFFRACTION

Diffractive x-section suppressed relative to Regge prediction as √s increases





Factor of ~8 (~5) suppression at √s = 1800 (540) GeV


RENORMALIZATION

Interpret flux as gap formation probability that saturates when it reaches unity

http://www.sciencedirect.com/science/article/pii/037026939501023J

Single Diffraction Renormalized - 1

KG ➔ CORFU-2001 http://arxiv.org/abs/hep-ph/0203141





2 independent variables: $t, \Delta y$

color factor

$$\kappa = \frac{g_{IP-IP-IP}(t)}{\beta_{IP-p-p}(0)} \approx 0.17$$

$$\frac{d^2\sigma}{dt\, d\Delta y} = C \cdot F_p^2(t) \cdot \left\{ e^{(\varepsilon + \alpha' t)\Delta y} \right\}^2 \cdot \kappa \cdot \left\{ \sigma_o\, e^{\varepsilon\, \Delta y'} \right\}$$

gap probability

sub-energy x-section

Gap probability ➔ (re)normalize it to unity

Single Diffraction Renormalized - 2

color factor

$$\kappa = \frac{g_{IP-IP-IP}(t)}{\beta_{IP-p-p}(0)} \approx 0.17$$

Experimentally ➔

$$\kappa = \frac{g_{IP-IP-IP}}{\beta_{IP-p}} = 0.17 \pm 0.02, \qquad \varepsilon = 0.104$$

KG&JM, PRD 59 (114017) 1999
http://dx.doi.org/10.1103/PhysRevD.59.114017

QCD: $\kappa = f_g \times \frac{1}{N_c^2 - 1} + f_q \times \frac{1}{N_c} \xrightarrow{Q^2 = 1} \approx 0.75 \times \frac{1}{8} + 0.25 \times \frac{1}{3} = 0.18$

Single Diffraction Renormalized - 3

$$\frac{d^2\sigma_{sd}(s,M^2,t)}{dM^2dt} = \left[\frac{\sigma_\circ}{16\pi}\sigma_\circ^{I\!Pp}\right] \frac{s^{2\epsilon}}{N(s,s_o)} \frac{e^{bt}}{(M^2)^{1+\epsilon}}$$

$$b = b_0 + 2\alpha' \ln\frac{s}{M^2}$$

$$s_o^{\rm CMG} = (3.7 \pm 1.5)\ {\rm GeV}^2$$

$$N(s,s_o) \equiv \int_{\xi_{\rm min}}^{\xi_{\rm max}} d\xi \int_{t=0}^{-\infty} dt\, f_{I\!P/p}(\xi,t) \overset{s\to\infty}{\to} \sim s_o^\epsilon \frac{s^{2\epsilon}}{\ln s}$$

←affects only the s-dependence

$$\frac{d^2\sigma_{sd}(s,M^2,t)}{dM^2dt} \overset{s\to\infty}{\to} \sim \ln s\, \frac{e^{bt}}{(M^2)^{1+\epsilon}}$$

set $N(s,s_o)$ to unity
➔ determines s_o

$$\sigma_{sd} \xrightarrow{s\to\infty} \sim \frac{\ln s}{b \to \ln s} \Rightarrow const$$

M^2 - Distribution: Data

➔ $d\sigma/dM^2|_{t=-0.05}$ ~ independent of s over 6 orders of magnitude!


http://physics.rockefeller.edu/publications.html

KG&JM, PRD 59 (1999) 114017

http://dx.doi.org/10.1103/PhysRevD.59.114017

Regge → data

$$\frac{d\sigma}{dM^2} \propto \frac{s^{2\varepsilon} \to 1}{(M^2)^{1+\varepsilon}}$$




➔ factorization breaks down to ensure M^2-scaling

Scale s_0 and *PPP* Coupling

Pomeron flux: interpreted as gap probability

➔set to unity: determines g_{PPP} and s_0

KG, PLB 358 (1995) 379 http://www.sciencedirect.com/science/article/pii/037026939501023J



Pomeron-proton x-section

- Two free parameters: s_0 and g_{PPP}
- Obtain product $g_{PPP}\, s_0^{\varepsilon/2}$ from σ_{SD}
- Renormalize Pomeron flux: determines s_o
- Get unique solution for g_{PPP}

DD at CDF

http://physics.rockefeller.edu/publications.html
http://dx.doi.org/10.1103/PhysRevLett.87.141802


$$\frac{d^3\sigma_{DD}}{dtdM_1^2dM_2^2} = \frac{d^2\sigma_{SD}}{dtdM_1^2}\frac{d^2\sigma_{SD}}{dtdM_2^2} \Big/ \frac{d\sigma_{el}}{dt}$$


Regge factorization


$$= \frac{[\kappa\beta_1(0)\beta_2(0)]^2}{16\pi} \frac{s^{2[\alpha(0)-1]}\, e^{b_{DD}t}}{(M_1^2M_2^2)^{1+2[\alpha(0)-1]}}$$

$$\frac{d^3\sigma_{DD}}{dtd\Delta\eta d\eta_c} = \left[\frac{\kappa\beta^2(0)}{16\pi} e^{2[\alpha(t)-1]\Delta\eta}\right]\left[\kappa\beta^2(0)\left(\frac{s'}{s_0}\right)^{\epsilon}\right]$$

gap probability x-section








←Regge

Regge ← RENORM

x-section divided by integrated gap prob.


SDD at CDF



http://physics.rockefeller.edu/publications.html

http://journals.aps.org/prl/abstract/10.1103/PhysRevLett.91.011802


- Excellent agreement between data and MBR (MinBiasRockefeller) MC




$$\frac{d^5\sigma}{dt_{\bar{p}}dtd\xi_{\bar{p}}d\Delta\eta d\eta_c} = \left[\frac{\beta(t)}{4\sqrt{\pi}}\, e^{[\alpha(t_{\bar{p}})-1]\ln(1/\xi)}\right]^2 \times \kappa\left\{\kappa\left[\frac{\beta(0)}{4\sqrt{\pi}} e^{[\alpha(t)-1]\Delta\eta}\right]^2 \kappa\left[\beta^2(0)\left(\frac{s''}{s_0}\right)^{\epsilon}\right]\right\}$$

CD/DPE at CDF

http://journals.aps.org/prl/abstract/10.1103/PhysRevLett.91.011802





- Excellent agreement between data and MBR-based MC

➔ Confirmation that both **low and high mass x-sections** are correctly implemented

RENORM Difractive Cross Sections

MBR MC Simulation in PYTHIA8➔http://arxiv.org/abs/1205.1446

$$\frac{d^2\sigma_{SD}}{dtd\Delta y} = \frac{1}{N_{\rm gap}(s)}\left[\frac{\beta^2(t)}{16\pi}e^{2[\alpha(t)-1]\Delta y}\right]\cdot\left\{\kappa\beta^2(0)\left(\frac{s'}{s_0}\right)^{\epsilon}\right\},$$

$$\frac{d^3\sigma_{DD}}{dtd\Delta ydy_0} = \frac{1}{N_{\rm gap}(s)}\left[\frac{\kappa\beta^2(0)}{16\pi}e^{2[\alpha(t)-1]\Delta y}\right]\cdot\left\{\kappa\beta^2(0)\left(\frac{s'}{s_0}\right)^{\epsilon}\right\},$$

$$\frac{d^4\sigma_{DPE}}{dt_1dt_2d\Delta ydy_c} = \frac{1}{N_{\rm gap}(s)}\left[\Pi_i\left[\frac{\beta^2(t_i)}{16\pi}e^{2[\alpha(t_i)-1]\Delta y_i}\right]\right]\cdot\kappa\left\{\kappa\beta^2(0)\left(\frac{s'}{s_0}\right)^{\epsilon}\right\}$$

$$\beta^2(t) = \beta^2(0)F^2(t)$$

$$F^2(t) = \left[\frac{4m_p^2-2.8t}{4m_p^2-t}\left(\frac{1}{1-\frac{t}{0.71}}\right)^2\right]^2 \approx a_1e^{b_1t}+a_2e^{b_2t}$$

α_1=0.9, α_2=0.1, b_1=4.6 GeV^{-2}, b_2=0.6 GeV^{-2}, s′=s $e^{-\Delta y}$, κ=0.17, $\kappa\beta^2(0)=\sigma_0$, s_0(units)=1GeV^2, σ_0=2.82 mb or7.25 GeV^{-2}

Total, Elastic, and Total Inelastic x-Sections

$$\sigma_{\rm ND} = (\sigma_{\rm tot} - \sigma_{\rm el}) - (2\sigma_{\rm SD} + \sigma_{\rm DD} + \sigma_{\rm CD})$$

CMG➔ R.J.M. Covolan [1], J. Montanha [2], K. Goulianos [3]
The Rockefeller University, 1230 York Avenue, New York, NY 10021, USA PLB 389, 196 (1996)

http://www.sciencedirect.com/science/article/pii/S0370269396013627

$$\sigma_{\rm tot}^{p^{\pm}p} = \begin{cases} 16.79 s^{0.104} + 60.81 s^{-0.32} \mp 31.68 s^{-0.54} & \text{for } \sqrt{s} < 1.8 \\ \sigma_{\rm tot}^{\rm CDF} + \frac{\pi}{s_0}\left[\left(\ln\frac{s}{s_F}\right)^2 - \left(\ln\frac{s^{\rm CDF}}{s_F}\right)^2\right] & \text{for } \sqrt{s} \geq 1.8 \end{cases}$$


KG MORIOND-2011 http://moriond.in2p3.fr/QCD/2011/proceedings/goulianos.pdf

$$\sqrt{s^{\rm CDF}} = 1.8 \text{ TeV}, \ \sigma_{\rm tot}^{\rm CDF} = 80.03 \pm 2.24 \text{ mb}$$


$$\sqrt{s_F} = 22 \text{ GeV} \qquad s_0 = 3.7 \pm 1.5 \text{ GeV}^2$$

$\sigma_{el}^{p\pm p} = \sigma_{tot}^{p\pm p} \times (\sigma_{el}/\sigma_{tot})^{p\pm p}$, with σ_{el}/σ_{tot} from CMG

- small extrapolation from 1.8 to 7 and up to 50 TeV



2009



$\sqrt{s_F}$=22 GeV

- Use the Froissart formula as a *saturated* cross section

$$\sigma_t(s > s_F) = \sigma_t(s_F) + \frac{\pi}{m^2} \cdot \ln^2 \frac{s}{s_F}$$

- This formula should be valid above the *knee* in σ_{sd} vs. $\sqrt{s}$ at $\sqrt{s_F} = 22$ GeV therefore valid at $\sqrt{s} = 1800$ GeV.

- Use $m^2 = s_o$ in the Froissart formula multiplied by $1/0.389$ to convert it to mb^{-1}.

- Note that contributions from Reggeon exchanges at $\sqrt{s} = 1800$ GeV are negligible, as can be verified from the global fit of CGM.

- Obtain the total cross section at the LHC:

$$\sigma_t^{\text{LHC}} = \sigma_t^{\text{CDF}} + \frac{\pi}{s_o} \cdot \left(\ln^2 \frac{s^{\text{LHC}}}{s_F} - \ln^2 \frac{s^{\text{CDF}}}{s_F} \right)$$

98 ± 8 mb at 7 TeV
109 ±12 mb at 14 TeV


Uncertainty is due to s_o

2015


Reduce Uncertainty in s_0

http://workshops.ift.uam-csic.es/LHCFPWG2015/program

EDS 2015: http://dx.doi.org/10.5506/APhysPolBSupp.8.783



Review of CEP by Albrow, Coughlin, Forshaw http://arxiv.org/abs/1006.1289
Fig from **Axial Field Spectrometer at** the CERN Intersecting Storage Rings



Data: Peter C. Cesil, AFS thesis
(courtesy Mike Albrow)
➔analysis: S and D waves
Conjecture: **tensor glue ball** (spin 2)
Fit: Gaussian

- $\langle M_{tgb} \rangle = \sqrt{s_0} = 2.10 \pm 0.68$ GeV
  - ➔$s_0 = 4.42 \pm 0.34$ GeV^2

20% increase in s_0
➔x-sections decrease

Predictions vs Measurements with/reduced Uncertainty in s_0

From my Moriond-2016 Talk

$\sqrt{s}$	MBR/Exp	σ_{tot}	σ_{el}	σ_{inel}
7 TeV	MBR	95.4±1.2	26.4±0.3	69.0±1.0
	TOTEM totem-lumInd	98.3±0.2±2.8 98.0±2.5	24.8±0.2±1.2 25.2±1.1	73.7±3.4 72.9±1.5
	ATLAS	95.35±1.36	24.00±0.60	71.34±0.90
8 TeV	MBR	97.1±1.4	27.2±0.4	69.9±1.0
	TOTEM	101.7±2.9	27.1±1.4	74.7±1.7
13 TeV	MBR	103.7±1.9	30.2±0.8	73.5±1.3
	ATLAS			σ_{inel}=73.1±0.9(exp)±6.6(lumi) ±3.8(extra.)mb

- RENORM/MBR with a **tensor-Pomeron model** predicts measured cross sections to the ~1% level
- **Test of RENORM/MBR**: ATLAS results using the ALFA and RP detectors to measure the cross sections

Stay tuned!

Totem 7 TeV http://arxiv.org/abs/1204.5689

Totem-Lum-Ind 7 TeV http://iopscience.iop.org/article/10.1209/0295-5075/101/21004

Atlas 7 TeV: http://arxiv.org/abs/1408.5778

Totem 8 TeV http://dx.doi.org/10.1103/PhysRevLett.111.012001

Atlas13 TeV Aspen 2016 Doug Schafer https://indico.cern.ch/event/473000/timetable/#all.detailed

Atlas/Totem 13TeV DIS15 https://indico.desy.de/contributionDisplay.py?contribId=330&confId=12482

Predictions vs Measurements w/reduced Uncertainty in s_o #1

ICNFP 2016

Slide from my ICNFP-2016 Talk

$\sqrt{s}$	MBR/Exp	Reference ➔next slide	S_{tot}	S_{el}	S_{inel}
7 TeV	MBR		95.4±1.2	26.4±0.3	69.0±1.0
	ATLAS	1	95.35±1.36	24.00±0.60	71.34±0.90
	TOTEM	2	101.7±2.9	27.1±1.4	74.7±1.7
	TOTEM_Lum_Ind	3	98.0±2.5	24.00±0.60	72.9±1.5
8 TeV	MBR		97.1±1.4	27.2±0.4	69.9±1.0
	TOTEM	4	101.7±2.9	27.1±1.4	74.7±1.7
13 TeV	MBR		103.7±1.9	30.2±0.8	73.5±1.3
	ATLAS	5 & 6		▫	73.1±0.9 (exp) ±6.6 (lumi) ±3.8 (extr)
	CMS	7			**71.3±0.5 (exp) ±2.1 (lumi) ±2.7 (extr)**

CONT➔

Predictions vs Measurements w/reduced Uncertainty in s_o #2

Caveat (slide from my ICNFP-2016 talk)

The MBR σ_{el} is larger than the ATLAS and the TOTEM_lum_Ind measurements by ~2 mb at $\sqrt{s}$=7 TeV, which might imply a higher MBR prediction at $\sqrt{s}$=13 TeV by 2-3 mb. Lowering the MBR σ_{el} prediction would lead to a larger σ_{inel}.This interplay between σ_{el} and σ_{inel} should be kept in mind as more results of σ_{el} and σ_{tot} at $\sqrt{s}$ = 13 TeV become available.

- RENORM/MBR with a **tensor-Pomeron model** predicts measured cross sections to the ~1% level
- **Test of RENORM/MBR**: ATLAS results using the ALFA and RP detectors to measure the cross sections

Stay tuned!

1) Atlas 7 TeV: http://arxiv.org/abs/1408.5778

2) Totem 7 TeV http://arxiv.org/abs/1204.5689

3) Totem-Lum-Ind 7 TeV http://iopscience.iop.org/article/10.1209/0295-5075/101/21004

4) Totem 8 TeV http://dx.doi.org/10.1103/PhysRevLett.111.012001

5) Atlas13 TeV Aspen 2016 D. Schafer https://indico.cern.ch/event/473000/timetable/#all.detailed

6) Atlas 13TeV DIS-2016 M. Trzebinski ttps://indico.desy.de/contributionDisplay.py?contribId=330&confId=12482

7) CMS 13TeV DIS-2016 H. Van Haevermaet https://indico.desy.de/contributionDisplay.py?contribId=105&confId=12482

MBR vs. ICHEP 2016 cross-section results

$\sqrt{s}$	MBR/Exp	Ref. # ➔slide 19	σ_{tot}	σ_{el}	σ_{inel}
7 TeV	MBR		95.4±1.2	26.4±0.3	69.0±1.0
	ATLAS	1	95.35±1.36	24.00±0.60	71.34±0.90
	TOTEM	2	101.7±2.9	27.1±1.4	74.7±1.7
	TOTEM_Lum_Ind	3	98.0±2.5	24.00±0.60	72.9±1.5
8TeV	MBR		97.1±1.4	27.2±0.4	69.9±1.0
				➔ATLAS vs. MBR in excellent agreement at 8 TeV	
	TOTEM ATLAS-ALFA fit	4 ICHEP16	101.7±2.9 **96.1±0.9**	27.1±1.4 24.3±0.4	74.7±1.7
13 TeV	MBR		103.7±1.9	30.2±0.8	73.5±1.3
	ATLAS ALFA-fit-result	5 & 6 ICHEP16			73.1±0.9 (exp) ±6.6 (lumi) ±3.8 (extr) **79.3±0.6(exp) ±1.3(lumi) ±2.5(extr)**
	CMS	7+ICHEP16			**71.3±0.5 (exp) ±2.1 (lumi) ±2.7 (extr)**

✓ Tomáš Sýkora, ICHEP16 x-sections summary talk http://ichep2016.org/

❑ At 13 TeV MBR is happy between the ATLAS and CMS ICHEP results
➔ awaiting settlement between the two experiments – keep tuned!

MBR vs. ICHEP 2016 cross-sections

$\sqrt{s}$ (TeV)	Input source	Reference*	σ_{tot} (mb)	σ_{el} (mb)	σ_{inel} (mb)
7	MBR	a	95.4 ± 1.2	26.4 ± 0.3	69.0 ± 1.0
	ATLAS	b	95.35 ± 1.36	24.00 ± 0.60	71.34 ± 0.90
	TOTEM	c	101.7 ± 1.36	27.1 ± 1.4	74.7 ± 1.7
	TOTEM_Lum_ind	d	98.0 ± 2.5	24.00 ± 0.60	72.9 ± 1.5
8	MBR	a	97.1 ± 1.4	27.2 ± 0.4	69.9 ± 1.0
	TOTEM	e	101.7 ± 2.9	27.1 ± 1.4	74.8 ± 1.7
	ATLAS_ALFA_fit	(h) ICHEP16	96.1 ± 0.9	24.3 ± 0.4	xxx
13	MBR	a	103.7 ± 1.9	30.2 ± 0.8	73.5 ± 1.3
	ATLAS	f&g	xxx	xxx	73.1 ± 0.9(exp)±3.8(extr)±6.6(lumi)
	ATLAS_ALFA_fit	(h) ICHEP16	xxx	xxx	79.3 ± 0.6(exp)±2.5(extr)±1.3(lumi)
	CMS	(h) ICHEP16	xxx	xxx	71.3 ± 0.6(exp)±2.7(extr)±0.1(lumi)

*Reference:
(a) http://arxiv.org/abs/1205.1446
(b) http://arxiv.org/abs/1408.5778
(c) http://arxiv.org/abs/1204.5689
(d) http://iopscience.iop.org/article/10.1209/0295-5075/101/21004
(e) http://dx.doi.org/10.1103/PhysRevLett.111.012001
(f) M. Trzebinski (ATLAS), DIS-2016 [7]-(a)
(g) H. Van Haevermaet (CMS), DIS-2016 [7]-(b)
(h) T. Sykora, *Cross sections summary*, ICHEP16 [8]

DIS-2017: MBR vs. TOTEM @ 2.76 TeV

https://indico.cern.ch/event/568360/ (from talk by Frigyes Nemes, slide #20)





TOTEM

σ_{tot} [mb]	σ_{el} [mb]	σ_{inel} [mb]
84.7 ± 3.3	21.8 ± 1.4	62.8 ± 2.9

MBR➔ 85.2 21.7 63.5

Syst. Uncertainty ~1.5% due to that in s_0

- Excellent agreement between TOTEM and MBR at 2.76 TeV
- Awaiting forthcoming results at 13 TeV from ATLAS, CMS, TOTEM

LHCC-2017: MBR vs. TOTEM @ 13 TeV

https://indico.cern.ch/event/679087/ (from talk by K. Osterberg)



$\sigma_{tot} = 110.6 \pm 3.4$ mb, $\sigma_{inel} = 79.5 \pm 1.8$ mb, $\sigma_{el} = 31.0 \pm 1.7$ mb



103.7±1.9	73.5±1.3	30.2±0.8

TOTEM paper➔
CERN-EP-2017-321
10 December 2017


Physics quantity	Value		Total uncertainty
	$\rho = 0.14$	$\rho = 0.1$	
B [GeV^{-2}]	20.36		$5.3 \cdot 10^{-2} \oplus 0.18 = 0.19$
σ_{tot} [mb]	109.5	110.6	3.4
σ_{el} [mb]	30.7	31.0	1.7
σ_{inel} [mb]	78.8	79.5	1.8
$\sigma_{el}/\sigma_{inel}$	0.390		0.017
σ_{el}/σ_{tot}	0.281		0.009

- Reasonable agreement between TOTEM and MBR predictions
- Possible Odderon effects not bincluded in MBR

Pythia8-MBR Hadronization Tune

An example of the diffractive tuning of PYTHIA-8 to the RENORM-NBR model

$$n_{ave} = \frac{\sigma_{QCD}}{\sigma_{IPp}}$$




Red line: best fit to multiplicity distributions.
(in bins of Mx, fits to higher tails only, default pT spectra)

R. Ciesielski, "Status of diffractive models", CTEQ Workshop 2013


SD and DD x-Sections vs Models

http://journals.aps.org/prd/abstract/10.1103/PhysRevD.92.012003




Single Diffraction

Double Diffraction



Monte Carlo Algorithm - Nesting



SUMMARY

- Review of RENORM predictions of diffractive physics
  - basic processes: SD1,SD2, DD, CD (DPE)
    - combined processes: multigap x-sections
    - ND ➔ no diffractive gaps: the only final state to be tuned
    - Monte Carlo strategy for the LHC – "nesting"
- Precision RENORM σ_{tot} prediction ^{W}/tensor glue-ball model
- ICHEP 2016
  - At 8 TeV ATLAS and MBR in excellent agreement
  - Disagreement betweenTOTEM and MBR persists
  - At 13 TeV MBR lies comfortably (!) between the ATLAS and CMS
- **LHCC-201: NEW➔TOTEM RESULTS at 8 and 13 TeV vs. MBR**
  - Agreement at 8 TeV, compatibility at 13 TeV
- **NESTING in MC simulation**

Thank you for your attention!